

Michael P. Carroll, MD, FAAOS, FACS
 Douglas E. Barkin, MD, FAAOS, FACS
 Robert J. Lippe, MD, FAAOS, FACS
 Jonathan B. Ticker, MD, FAAOS
 Armand E. Abulencia, MD
 James J. Morimando, PA-C
 Karen A. Andrews, PA-C
 Kevin M. Condon - Administrator



Arthroscopy & Sports Medicine
 Hip & Knee Replacement
 Shoulder Surgery
 Knee Surgery
 Spine Surgery
 Fracture Care
 Hand & Elbow Surgery
 Ankle & Foot Disorders

RADIOGRAPHS FOR SHOULDER SERIES

(Please perform those radiographs indicated in the boxes checked.)

1. True AP Shoulder (Glenohumeral Joint) View
 A. Patient position: erect, affected side against Bucky, turn patient 30-35°
 B. Tube: C.R. perpendicular to patient
 C. Distance: 48"
 D. Settings (average male): 15-20 MAS, 75 KVP, regular cassette
2. Outlet ("Y") View
 A. Patient position: erect PA, affected side against Bucky, turn patient 40-45°
 B. Tube: C.R. 15-20° caudad at coracoid (along spine of scapula)
 C. Distance: 48"
 D. Settings (average male): 15-20 MAS, 85 KVP, regular cassette
3. Axillary View
 A. Patient position: supine with small sponge under shoulder, humerus 90° to body (if possible), elbow bent 90° and perpendicular to table
 B. Tube: C.R. bisecting angle of humerus and body
 C. Distance: 40"
 D. Settings (average male): 7 MAS, 62 KVP, regular cassette
4. Zanca (Acromioclavicular Joint) View
 A. Patient position: standing in AP position
 B. Tube: C.R. 5-15° cephalad bisecting the AC joint
 C. Distance: 40"
 D. Settings (average male): 15-20 MAS, 62 KVP, regular cassette
5. Stryker Notch View
 A. Patient position: supine, cassette posterior to shoulder, with humerus flexed approximately 120° so that the hand can be placed on top of patient's head
 B. Tube: C.R. 10° cephalad centered over coracoid process
 C. Distance: 40"
 D. Settings (average male): 20 MAS, 75-80 KVP, regular cassette Bucky
6. Velpeau Axillary View
 A. Patient position: standing or sitting at edge of table leaning back 20-30° over table; cassette on table beneath shoulder
 B. Tube: C.R. passes vertically from superior to inferior, through shoulder joint
 C. Distance: 48"
 D. Settings (average male): 20 MAS, 85 KVP, regular cassette

